

# Pakistan soil transmitted helminth survey data

---

## Mapped references

1. Mehraj V, Hatcher J, Akhtar S, Rafique G, Beg MA (2008) Prevalence and factors associated with intestinal parasitic infection among children in an urban slum of Karachi. *PLoS One* 3: e3680
2. Ensink JH, van der Hoek W, Mukhtar M, Tahir Z, Amerasinghe FP (2005) High risk of hookworm infection among wastewater farmers in Pakistan. *Trans R Soc Trop Med Hyg* 99: 809-818
3. Chaudhry ZH, Afzal M, Malik MA (2004) Epidemiological factors affecting prevalence of intestinal parasites in children of Muzaffarabad District. . *Pakistan Journal of Zoology* 36: 267-271
4. Ahmed AK, Malik B, Shaheen B, Yasmeen G, Dar JB, Mona AK, Gulab S, Ayub M (2003) Frequency of intestinal parasitic infestation in children of 5-12 years of age in Abbottabad. *J Ayub Med Coll Abbottabad* 15: 28-30
5. Nishiura H, Imai H, Nakao H, Tsukino H, Changazi MA, Hussain GA, Kuroda Y, Katoh T (2002) Ascaris lumbricoides among children in rural communities in the Northern Area, Pakistan: prevalence, intensity, and associated socio-cultural and behavioral risk factors. *Acta Trop* 83: 223-231

## Unmapped references

1. Feenstra S (2000) Unpublished data contributed by author, Bahalwanagar, Punjab